

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07D 309/30, A61K 31/36	A1	(11) International Publication Number: WO 94/29292 (43) International Publication Date: 22 December 1994 (22.12.94)
(21) International Application Number: PCT/SI94/00010 (22) International Filing Date: 8 June 1994 (08.06.94) (30) Priority Data: P 9300303 8 June 1993 (08.06.93) SI (71) Applicant (for all designated States except US): KRKA TOVARNA ZDRAVIL P.O. [SI/SI]; Cesta herojev 45, 68000 Novo mesto (SI). (72) Inventors; and (75) Inventors/Applicants (for US only): HAJKO, Pavica [SI/SI]; Majde Šilc 19, 68000 Novo mesto (SI). VESEL, Tanja [SI/SI]; V brezov Iog 5, 68000 Novo mesto (SI). RADEŽ, Ivan [SI/SI]; Brš 25, 68000 Novo mesto (SI). POKORNY, Miroslav [SI/SI]; Cankarjeva 11, 68000 Novo mesto (SI). (74) Agent: PATENTNA PISARNA D.O.O.; Čopova 14, 61000 Ljubljana (SI).		(81) Designated States: BG, BY, CA, CZ, FI, GE, HU, KG, LV, MD, NO, PL, RO, RU, SK, TJ, UA, US, UZ, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: PROCESS FOR THE ISOLATION OF LOVASTATIN		
(57) Abstract <p>There is disclosed a process for the isolation of the hypolipaeic active substance lovastatin from a sample, e.g. fermentation broth, mycelium or filtrate of <i>Aspergillus terreus</i> (ATCC 20542) or <i>Aspergillus oryzae</i> (ATCC 74135) by extraction with butyl acetate. Simultaneously with the concentration of the extract, lactonisation takes place. There follows a direct crystallization of lovastatin in the lactone form from butyl acetate.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.


AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

PROCESS FOR THE ISOLATION OF LOVASTATIN

Technical Field


The invention belongs to the field of biotechnology and relates to a process for the isolation of the hypolipaeic active substance lovastatin from a fermentation broth. Lovastatin is produced as a secondary metabolite of the fungus *Aspergillus terreus* (US 4,231,938) deposited in American Type Culture Collection under Nos. ATCC 20541, ATCC 20542, and *Monascus ruber* deposited in Fermentation Research Institute Agency of Industrial Science and Technology, Ministry of International Trade and Industry, Japan (DE 30 06 216 A1) under No. Ferm 4822. Other kinds of microorganisms producing lovastatin are known as well, e.g. a mutant of the microorganism *Aspergillus terreus* and *Aspergillus oryzae* marked ATCC 74135.

Lovastatin is chemically 1',2',6',7',8a'-hexahydro-3,5-dihydroxy-2',6'-dimethyl-8'-2''-methyl-1''-oxobutoxy)-1-naphtalene heptanoic acid-5-lactone (Stubbs et al., 1986) of the formula (EP 0 033 537 A1)



Compound I

An active form of lovastatin is also an acid, which is chemically 1,2,6,8,8a-hexahydro- β,δ -dihydroxy-1-naphtalene heptanoic acid (Alberts et al., 1980) of the formula (EP 0 022 478 A1)



Compound II

The lactone form of lovastatin is used as an agent for reducing cholesterol level in blood (Scott M.G. and Vega G.L., 1985). It inhibits the biosynthesis of mevalonic acid by inhibition of 3-hydroxy-3-methylglutaryl A reductase coenzyme (HMG-CoA reductase, E.C. 1.1.1.34) (Zubay et al., 1984).

Prior Art

After the completed fermentation, lovastatin is present in the broth in the lactone form (compound I) and in the acid form (compound II). In the isolation process as disclosed in EP 0 033 536 A2, lovastatin is extracted from the broth with ethyl acetate. The extract is concentrated by vacuum distillation. Since lovastatin is present in the lactone form as well as in the acid form and only the lactone is of commercial interest, the acid form should be converted into the lactone. The lactonisation is carried out by the reflux of the concentrate in toluene at 106°C for 2 hours. After the lactonisation is complete, the solution is concentrated to a small volume. A pure substance is obtained by means of purifying the concentrate on columns packed with silica gel, in the presence of solvents such as ethyl acetate or n-hexane. The collected fractions are again concentrated *in vacuo* and then pure lovastatin crystallizes in the lactone form.

Due to the sophisticated multi-step procedure and vigorous conditions applied during the isolation, the yields of lovastatin are generally low. Different solvents, which in part exhibit toxicity, are used such as benzene, toluene, acetonitrile or ethyl acetate. Hence working with these solvents endangers the health of the persons involved and poses a problem with respect to the environment.

Description of the Invention

An object of the present invention was to overcome the above deficiencies of the state of the art and to provide a process for isolating lovastatin in an efficient and economic way. In addition, attention should be paid to health aspects of the workers involved in the isolation and to the environment.

As a result of extensive search leading to the present invention, the present inventors have found that lovastatin may be easily isolated by extracting lovastatin from a sample with butyl acetate, concentrating the solution and carrying out a crystallization.

In a preferred embodiment the concentration of the solution is carried out under reduced pressure. In a further preferred embodiment concentration is carried out at a temperature above 40 °C with or without reduced pressure.

In yet another preferred embodiment the sample of lovastatin is derived from microorganisms, preferably a fungus capable of producing lovastatin. In this case the sample may be the fermentation broth in which the microorganism or the fungus, respectively, has been grown, the mycelium of the fungus or the filtrate.

The fungus to be used is not limited and any fungus capable of producing lovastatin may be utilized, with those belonging to the genus *Aspergillus*, preferably *Aspergillus oryzae* or *Aspergillus terreus*, being preferred.

In yet another preferred embodiment the fungus is *Aspergillus terreus* (ATCC 20542) or *Aspergillus oryzae* (ATCC 74135).

According to the process of the present invention lovastatin may be obtained exclusively in the lactone form. It has surprisingly been found that the solvent utilized, i.e. butyl acetate, provides features enabling to achieve this goal. Namely, in the

lactonisation process water is released and should be removed from the reaction medium. Since butyl acetate has a higher boiling point and does not substantially form azeotropic mixtures with water under the given conditions, essentially all water formed during the lactonisation process may be removed.

By the process according to the present invention it has become possible to substantially shorten the time for isolating lovastatin. Since butyl acetate may be used to extract lovastatin from a sample and at the same time may be utilized for performing the lactonisation, the toxic effects of other solvents may be obviated. The process may be industrially reproduced with the final yields of about 60%, irrespective of lovastatin being produced by microorganisms intracellularly or extracellularly, in the lactone form or in the acid form.

The isolated substance is the same as the substance isolated according to Prior Art processes. The structure was confirmed by IR spectroscopy (Fig.1), mass spectroscopy (Fig. 2), NMR (Fig. 3) and UV spectroscopy (Fig. 4).

The invention is illustrated by the following non-limiting Examples.

Example 1

Fermentation broth (1000 ml) obtained by the fermentation of *Aspergillus terreus* (ATCC 20542) or *Aspergillus oryzae* (ATCC 74135) and having a pH value of 4.8, an age of 120 hours and a lovastatin content of 300 µg/ml (lactone + acid) was acidified with HCl (36%) to a pH of 3-5. The fermentation broth was then cooled below 20 °C, butyl acetate (500 ml) was added and it was extracted for 4 hours at this temperature. After the extraction was complete, it was centrifuged and the aqueous and organic phases were separated. The mycelium and the aqueous phase were discarded and the organic phase was further treated. It was concentrated by vacuum distillation at a temperature above 40 °C to a volume of 50 ml. Simultaneously with the concentration, the lactonisation process took place as well. The concentrate was then cooled to a temperature below 20 °C and left to stand a few hours so that lovastatin crystallized. The obtained crude product was recrystallized. Lovastatin (170 mg) in the lactone form having a purity above 90% was obtained.

Example 2

Fermentation broth (1000 ml) obtained by the fermentation of *Aspergillus terreus* (ATCC 20542) or *Aspergillus oryzae* (ATCC 74135) and having a pH value of 4.73, an age of 120 hours and a lovastatin content of 190 µg/ml (lactone + acid) in the filtrate and 15 µg/ml in the mycelium was acidified with HCl (36%) to a pH of 3-5. The broth was filtered over Oliver filter. The mycelium was discarded, the filtrate was cooled to a temperature below 20 °C and half the volume of butyl acetate was added. It was extracted for 1 hour. Then it was proceeded as in Example 1. Lovastatin (79 mg) in the lactone form having a purity above 90 % was obtained.

Example 3

Fermentation broth (1000 ml) obtained by the fermentation of *Aspergillus terreus* (ATCC 20542) or *Aspergillus oryzae* (ATCC 74135) and having a pH value value of 4.78, an age of 120 hours and a lovastatin content of 5 µg/ml (lactone + acid) in the filtrate and 300 µg/ml in the mycelium was acidified with HCl (36%) to a pH of 3-5 and centrifuged. The filtrate was discarded, a 4-fold volume of butyl acetate was added to the mycelium and it was cooled below 20 °C. It was extracted for 4 hours. The extraction was followed by centrifuging. The mycelium was discarded and the filtrate treated as in Example 1. Lovastatin (90 mg) in the lactone form having a purity above 90 % was obtained.

CLAIMS

1. A process for the isolation of lovastatin, characterized in that lovastatin is extracted from a sample with butyl acetate, concentrated and left to crystallize.
2. A process according to Claim 1, wherein the solution of lovastatin in butyl acetate is concentrated under reduced pressure.
3. A process according to Claim 1 or 2, wherein the solution of lovastatin in butyl acetate is concentrated at a temperature above 40 °C.
4. A process according to any of the Claims 1 to 3, wherein lovastatin is extracted from the fermentation broth, mycelium or filtrate of a microorganism capable of producing lovastatin with butyl acetate.
5. A process according to claim 4, wherein the microorganism capable of producing lovastatin is a fungus.
6. A process according to claim 5, wherein the fungus belongs to the genus *Aspergillus*.
7. A process according to Claim 6, wherein the fungus belongs to the genus *Aspergillus terreus* or *Aspergillus oryzae*.
8. A process according to Claim 7, wherein the fungus is *Aspergillus terreus* (ATCC 20542) or *Aspergillus oryzae* (ATCC 74135).

1/4




FIG. 1




FIG. 2




FIG. 3

4/4




FIG. 4

INTERNATIONAL SEARCH REPORT

International Application No

PCT/SI 94/00010

A. CLASSIFICATION OF SUBJECT MATTER
IPC 5 C07D309/30 A61K31/36

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 5 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,A	WO,A,94 10328 (BIOGAL GYOGYSZERGYAR RT) 11 May 1994 see page 1, line 18 ---	1-8
P,A	EP,A,0 556 699 (NOVOPHARM LTD.) 25 August 1993 see page 4, line 54 ---	1-8
A	EP,A,0 033 536 (MERCK & CO. INC.) 12 August 1991 cited in the application see page 5, line 14 ---	1-8
A	EP,A,0 022 478 (MERCK & CO. INC.) 21 January 1981 cited in the application see page 14, line 22 ---	1-8
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

*** Special categories of cited documents :**

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

9 September 1994

Date of mailing of the international search report

20. 09. 94

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+ 31-70) 340-3016

Authorized officer

Gettins, M

INTERNATIONAL SEARCH REPORT

nal Application No

PCT/SI 94/00010

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DE,A,30 06 216 (SANKYO COMPANY LIMITED) 4 September 1980 cited in the application see page 8, line 19 - line 20 ---	1-8
A	US,A,4 231 938 (MONAGHAN) 4 November 1980 cited in the application see column 6, line 11 - line 12 -----	1-8

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/SI 94/00010

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9410328	11-05-94	NONE	
EP-A-0556699	25-08-93	AU-A- 3212193 BG-A- 97421 JP-A- 6022780 PL-A- 297691 CA-A- 2062023	12-08-93 24-03-94 01-02-94 24-01-94 11-08-93
EP-A-0033536	12-08-81	US-A- 4282155 AT-T- 9344 JP-B- 1038116 JP-C- 1553297 JP-A- 56122373	04-08-81 15-09-84 11-08-89 04-04-90 25-09-81
EP-A-0022478	21-01-81	US-A- 4231938 AT-T- 2620 AU-B- 535944 AU-A- 5901780 CA-A- 1161380 JP-C- 1518752 JP-A- 57163374 JP-B- 63066306 JP-C- 1319910 JP-A- 56008689 JP-B- 58016875 US-A- 4294926 US-A- 4319039 US-A- 4294846 US-A- 4342767	04-11-80 15-03-83 12-04-84 18-12-80 31-01-84 07-09-89 07-10-82 20-12-88 29-05-86 29-01-81 02-04-83 13-10-81 09-03-82 13-10-81 03-08-82
DE-A-3006216	04-09-80	JP-C- 1247674 JP-A- 55111790 JP-B- 59025599 AT-B- 373915 AU-A- 5567380 BE-A- 881825 CA-A- 1129794 CH-A- 645890 DE-A, C 3051175 FR-A, B 2449685	16-01-85 28-08-80 19-06-84 12-03-84 28-08-80 20-08-80 17-08-82 31-10-84 22-06-89 19-09-80

INTERNATIONAL SEARCH REPORT

information on patent family members

nal Application No

PCT/SI 94/00010

PCT/81 94/00010

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE-A-3006216		GB-A, B 2046737	19-11-80
		NL-A- 8001041	22-08-80
		SE-B- 453301	25-01-88
		SE-A- 8001339	21-08-80
		SU-A- 1158048	23-05-85
		SU-A- 969702	30-10-82

US-A-4231938	04-11-80	AT-T- 2620	15-03-83
		AU-B- 535944	12-04-84
		AU-A- 5901780	18-12-80
		CA-A- 1161380	31-01-84
		EP-A, B 0022478	21-01-81
		JP-C- 1518752	07-09-89
		JP-A- 57163374	07-10-82
		JP-B- 63066306	20-12-88
		JP-C- 1319910	29-05-86
		JP-A- 56008689	29-01-81
		JP-B- 58016875	02-04-83
		US-A- 4294926	13-10-81
		US-A- 4319039	09-03-82
